


Claims

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

R¹ represents hydrogen, -C₁₋₆ alkyl, -C₀₋₄ alkyl-C₃₋₈cycloalkyl, -C₂₋₄ alkoxy-C₁₋₄ alkyl, -C₁₋₄ alkyl-aryl, -C₁₋₄ alkyl-heteroaryl or -C₀₋₄ alkyl-heterocyclyl, or R¹ is linked to R² to form a group (CH₂)₂, (CH₂)₃ or (CH₂)₄;

wherein said alkyl, cycloalkyl, aryl, heteroaryl or heterocyclyl groups of R¹ may be optionally substituted by one or more (e.g. 1, 2 or 3) halogen, C₁₋₆ alkyl, C₁₋₆ alkoxy, cyano, amino or trifluoromethyl groups;

R² represents hydrogen or C₁₋₆ alkyl;

m represents an integer from 1 to 4, such that when m is an integer greater than 1, two R² groups may instead be linked to form a CH₂, (CH₂)₂ or (CH₂)₃ group;

R³, R⁴ and R⁵ independently represent hydrogen, halogen, cyano, -CF₃, -CF₃O, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkanoyl or a group -CONR⁶R⁷;

R⁶ and R⁷ independently represent hydrogen or C₁₋₆ alkyl or R⁶ and R⁷ together with the nitrogen to which they are attached may form a nitrogen containing heterocyclyl or heteroaryl group;

n represents an integer from 1 to 3;

p represents 1 or 2;

J represents CH₂, CO, CF₂, CH(OR⁸), NR⁹, SO, O or S;

R⁸ and R⁹ independently represent hydrogen or C₁₋₆ alkyl;

A represents an -aryl, -heteroaryl, -aryl-aryl, -aryl-heteroaryl, -heteroaryl-aryl or -heteroaryl-heteroaryl group;

wherein said aryl and heteroaryl groups of A may be optionally substituted by one or more (e.g. 1, 2 or 3) substituents which may be the same or different, and which are selected from the group consisting of halogen, hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, C₁₋₆ alkyl, trifluoromethanesulfonyloxy, pentafluoroethyl, C₁₋₆ alkoxy, arylC₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkoxyC₁₋₆ alkyl, C₃₋₇ cycloalkylC₁₋₆ alkoxy, C₁₋₆ alkanoyl, C₁₋₆ alkoxycarbonyl, C₁₋₆ alkylsulfonyl, C₁₋₆ alkylsulfinyl, C₁₋₆ alkylsulfonyloxy, C₁₋₆ alkylsulfonylC₁₋₆ alkyl, arylsulfonyl, arylsulfonyloxy, arylsulfonylC₁₋₆ alkyl, C₁₋₆

alkylsulfonamido, C₁₋₆ alkylamido, C₁₋₆ alkylsulfonamidoC₁₋₆ alkyl, C₁₋₆ alkylamidoC₁₋₆ alkyl, arylsulfonamido, arylcarboxamido, arylsulfonamidoC₁₋₆ alkyl, arylcarboxamidoC₁₋₆ alkyl, aroyl, aroylC₁₋₆ alkyl, arylC₁₋₆ alkanoyl, or a group CONR¹⁰R¹¹ or SO₂NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or C₁₋₆ alkyl or R¹⁰ and R¹¹

- 5 together with the nitrogen atom to which they are attached may form a nitrogen containing heterocyclyl or heteroaryl group;
or solvates thereof.
2. A compound of formula (I) as defined in claim 1 which is:
10 3-[(3-Chlorophenyl)methyl]-8-(1-piperazinyl)quinoline;
(3-Chlorophenyl)[8-(1-piperazinyl)-3-quinoliny]methanone;
3-(Phenyloxy)-8-(1-piperazinyl)quinoline;
or a pharmaceutically acceptable salt thereof.
- 15 3. A pharmaceutical composition which comprises a compound as defined in claim 1 or claim 2 and a pharmaceutically acceptable carrier or excipient.
4. A compound as defined in claim 1 or claim 2 for use in therapy.
- 20 5. A compound as defined in claim 1 or claim 2 for use in the treatment of depression, anxiety, Alzheimer's disease, age related cognitive decline, ADHD, obesity, mild cognitive impairment, schizophrenia, cognitive deficits in schizophrenia and stroke.
- 25 6. The use of a compound of formula (I) as defined in claim 1 or claim 2 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the treatment or prophylaxis of depression, anxiety, Alzheimer's disease, age related cognitive decline, ADHD, obesity, mild cognitive impairment, schizophrenia, cognitive deficits in schizophrenia and stroke.
- 30 7. A pharmaceutical composition comprising a compound of formula (I) as defined in claim 1 or claim 2 for use in the treatment of depression, anxiety, Alzheimer's disease, age related cognitive decline, ADHD, obesity, mild cognitive impairment, schizophrenia, cognitive deficits in schizophrenia and stroke.
- 35 8. A method of treating depression, anxiety, Alzheimer's disease, age related cognitive decline, ADHD, obesity, mild cognitive impairment, schizophrenia, cognitive deficits in schizophrenia and stroke which comprises administering a safe and therapeutically effective amount to a patient in need thereof of a compound of formula (I) as defined in claim 1 or claim 2 or a pharmaceutically acceptable salt thereof.